

ZnO nanorods for efficient third harmonic UV generation: erratum

Susanta Kumar Das,^{1,4} Frank Güell,² Ciarán Gray,³ Prasanta Kumar Das,⁴ Ruediger Grunwald,¹ and Enda McGlynn^{3,*}

¹Max-Born-Institut für Nichtlineare Optik und Kurzzeitspektroskopie, Max-Born-Strasse 2a, D-12489 Berlin, Germany

²Departament d'Electrònica, Universitat de Barcelona, C/Martí i Franquès 1, E-08028 Barcelona, Catalunya, Spain

³School of Physical Sciences, National Centre for Plasma Science and Technology, Dublin City University, Glasnevin, Dublin 9, Ireland

⁴School of Applied Sciences, KIIT University, Bhubaneswar - 751024, Odisha, India

*enda.mcglynn@dcu.ie

Abstract: We correct Eq. (1) of our previous paper showing the pulse duration (τ) dependence on group delay dispersion (β_2) and initial duration of the unchirped pulse coming from the laser (τ_0). The correct form of the equation was used in all the calculations in the previous paper however, so in all other respects our previous results and conclusions are unchanged.

©2014 Optical Society of America

OCIS codes: (160.4236) Nanomaterials; (190.4400) Nonlinear optics, materials; (190.7110) Ultrafast nonlinear optics.

References and links

1. S. K. Das, F. Guell, C. Gray, P. K. Das, R. Grunwald, and E. McGlynn, "ZnO nanorods for efficient third harmonic UV generation," *Opt. Mater. Express* **4**(4), 701–709 (2014).
2. D. Marcuse, "Pulse distortion in single-mode fibers," *Appl. Opt.* **19**(10), 1653–1660 (1980).
3. I. Walmsley, L. Waxer, and C. Dorrer, "The role of dispersion in ultrafast optics," *Rev. Sci. Instrum.* **72**(1), 1–29 (2001).

In Eq. (1) of article [1], we gave an equation showing the dependence of pulse duration (τ) on group delay dispersion (β_2) and initial duration of the unchirped pulse coming from the laser (τ_0). There are typographical errors in this equation in article [1] and the corrected equation is given below in Eq. (1), as may be verified in a number of other sources (such as [2] and [3]):

$$\tau = \tau_0 \sqrt{1 + \left(4 \cdot \ln(2) \cdot \left(\frac{\beta_2}{\tau_0^2} \right) \right)^2} \quad (1)$$

The correct form of the equation was used in all the calculations in the previous paper [1] however, so in all other respects our previous results and conclusions are unchanged.