

# Author Queries

*Journal:* Philosophical Transactions of the Royal Society B

*Manuscript:* RSTB20180359

As the publishing schedule is strict, please note that this might be the only stage at which you are able to thoroughly review your paper.

Please pay special attention to author names, affiliations and contact details, and figures, tables and their captions.

The corresponding author must provide an ORCID ID if they haven't done so already. If you or your co-authors have an ORCID ID please supply this with your corrections. More information about ORCID can be found at <http://orcid.org/>.

No changes can be made after publication.

**SQ1** Your supplementary material will be published online alongside your article and on [rs.figshare.com](http://rs.figshare.com) exactly as the file(s) are provided. Therefore, please could you either confirm that your supplementary material is correct, or – if you have any changes to make to these files – email these along with your proof corrections to the journal inbox. Your ESM files are listed here for your convenience:

RSTB-2018-0359\_revised\_SupplementalMaterial.pdf

RSTB-2018-0359\_Exp1\_RAWDATA.xlsx

RSTB-2018-0359\_Exp2\_RAWDATA.xlsx

**Q1** We have included section heading Methods for consistency with Section 3. Please check and confirm whether this is ok.

**Q2** Please update reference [43].

**Q3** While the online version of figure 3 will be in colour, we have been instructed to print the figure in black and white. Please note that if you have explicitly referred to colour in the caption this may affect the legibility of the figure in print.

## Research



**Cite this article:** O'Dowd A, Cooney SM, McGovern DP, Newell FN. 2019 Do synaesthesia and mental imagery tap into similar cross-modal processes? *Phil. Trans. R. Soc. B* 20180359.  
<http://dx.doi.org/10.1098/rstb.2018.0359>

Accepted: 2 June 2019

One contribution of 16 to a discussion meeting issue 'Bridging senses: new developments in synaesthesia'.

**Subject Areas:**

behaviour, cognition, neuroscience

**Keywords:**

synaesthesia, mental imagery, touch, somatosensation, mirror-touch, cross-modal correspondences

**Author for correspondence:**

Fiona N. Newell

e-mail: [fiona.newell@tcd.ie](mailto:fiona.newell@tcd.ie)

†Present address: Institute of Neuroscience, Lloyd Building, Trinity College Dublin, Dublin D02 PN40, Ireland.

Electronic supplementary material is available online at [rs.figshare.com](http://rs.figshare.com).

# Do synaesthesia and mental imagery tap into similar cross-modal processes?

Alan O'Dowd<sup>1</sup>, Sarah M. Cooney<sup>1</sup>, David P. McGovern<sup>1,2</sup> and Fiona N. Newell<sup>1,†</sup>

<sup>1</sup>School of Psychology and Institute of Neuroscience, Trinity College Dublin, Ireland

<sup>2</sup>School of Psychology, Dublin City University, Dublin, Ireland

FNN, 0000-0002-7363-2346

Synaesthesia has previously been linked with imagery abilities, although an understanding of a causal role for mental imagery in broader synaesthetic experiences remains elusive. This can be partly attributed to our relatively poor understanding of imagery in sensory domains beyond vision. Investigations into the neural and behavioural underpinnings of mental imagery have nevertheless identified an important role for imagery in perception, particularly in mediating cross-modal interactions. However, the phenomenology of synaesthesia gives rise to the assumption that associated cross-modal interactions may be encapsulated and specific to synaesthesia. As such, evidence for a link between imagery and perception may not generalize to synaesthesia. Here, we present results that challenge this idea: first, we found enhanced somatosensory imagery evoked by visual stimuli of body parts in mirror-touch synaesthetes, relative to other synaesthetes or controls. Moreover, this enhanced imagery generalized to tactile object properties not directly linked to their synaesthetic associations. Second, we report evidence that the concurrent evoked by grapheme–colour synaesthetes was sufficient to trigger visual-to-tactile correspondences that are common to all. Together, these findings show that enhanced mental imagery is a consistent hallmark of synaesthesia, and suggest the intriguing possibility that imagery may facilitate the cross-modal interactions that underpin synaesthetic experiences.

This article is part of the discussion meeting issue 'Bridging senses: new developments in synaesthesia'.

## 1. Introduction

Synaesthesia has long been compared with imagery [1], due to similarities between the phenomenological experiences [2] and that both experiences occur in the absence of an exogenous stimulus. There is compelling evidence that synaesthesia is associated with vivid mental imagery (e.g. [3,4] for reviews), particularly from studies of grapheme–colour synaesthesia [5,6] but also within other types, such as space–sequence synaesthesia ([4,7–10], although see [11]). The association between imagery abilities and synaesthesia seems to be quite nuanced, however. For example, imagery appears to be particularly enhanced within the modality that is specific to the inducer or concurrent [12], but is not constrained to the specific stimulus categories involved in the synaesthetic experience itself. Instead, enhanced visual imagery generalizes to other visual stimuli, at least in grapheme–colour synaesthetes [5].

Given that imagery affects perceptual [13,14] and memory [15] processes, and shares neural substrates underpinning perceptual function [16,17], a general enhancement of imagery abilities in synaesthetes may lead to better performance in perceptual tasks within the relevant modality. Indeed, grapheme–colour synaesthesia is associated with better colour perception [18], mirror-touch synaesthesia (MTS) with more sensitive tactile perception [19] and space–sequence synaesthesia with more efficient perception of rotated stimuli [8,9]. Studies that have formally investigated a link between imagery and synaesthesia often point to similarities between the two experiences

64 regarding their effects on perception. For example, Alvarez &  
 65 Robertson [20] found that when a congruent prime comprised  
 66 both a real colour and induced the same synaesthetic colour,  
 67 priming was enhanced in a colour naming task relative to  
 68 either colour type alone. Moreover, this congruency effect cor-  
 69 related with synaesthetes' reported vividness of imagery,  
 70 suggesting a strong connection between the two experiences.  
 71 Chiou *et al.* [21] conducted a colour priming task using bin-  
 72 ocular rivalry and reported that while both a synaesthetic  
 73 and imagined colour prime facilitated colour perception, a  
 74 real colour prime tended to suppress colour perception  
 75 when the prime and target stimulated the same monocular  
 76 location. An important question therefore arises about the  
 77 nature of the distinction between synaesthetic, imagined and  
 78 real perceptual experiences which, according to these studies,  
 79 may be influenced by the specific properties of the respective  
 80 stimulation (e.g. global or local) as well as the origin of the  
 81 phenomenon itself (endogenous or exogenous).

82 Apart from evidence that imagery is associated with  
 83 synaesthesia, and that both imagery and synaesthesia show  
 84 similar effects on perception, other studies have provided evi-  
 85 dence that imagery is sufficient to mediate synaesthesia.  
 86 Synaesthetic experiences themselves can be generated even  
 87 when the inducing stimulus is not physically present in the  
 88 environment, by being evoked through wilful imagery [6]  
 89 or even by concept formation (see [22,23]). However, to  
 90 date, efforts to demonstrate links between synaesthesia and  
 91 mental imagery abilities have mainly been confined to the  
 92 visual domain (e.g. colour or space-sequences). Indeed, Spil-  
 93 ler *et al.* [12] reported more vivid mental imagery in  
 94 synaesthetes than non-synaesthetes when the sensory  
 95 modality of the inducer and concurrent was congruent.  
 96 Thus, when both the inducer and concurrent are visual, it  
 97 might be possible that imagery is enhanced due to repeated  
 98 stimulation within the visual system. What is less clear is  
 99 whether imagery would also consistently be enhanced  
 100 across modalities when the synaesthetic concurrent arises in  
 101 a sensory modality that differs to that of the inducing stimu-  
 102 lus [12]. The nature and vividness of imagery in incidences  
 103 of synaesthesia that are triggered across modalities, and in  
 104 modalities other than vision, have rarely been investigated.

105 In typical perceptual processing, imagery can influence  
 106 multisensory perception (e.g. [24,25]). For example, Berger  
 107 & Ehrsson [26] reported that imagined sounds can trigger  
 108 similar cross-modal illusions as their real counterparts, includ-  
 109 ing the ventriloquist effect and McGurk illusion. Furthermore,  
 110 Lacey *et al.* [27] argued for a central role of imagery in the  
 111 interactions underpinning the cross-modal perception of  
 112 objects. It is therefore possible that imagery abilities in  
 113 synaesthesia may be linked to the cross-modal interactions  
 114 underpinning the experience and that, moreover, synaesthesia  
 115 (like imagery) may affect cross-modal interactions that are  
 116 commonly experienced by all.

117 One intriguing aspect of synaesthetic experiences is that  
 118 there appears to be little evidence that the concurrent affects  
 119 the perception of stimuli beyond the specific synaesthetic  
 120 association itself. In other words, there is evidence to suggest  
 121 that colour concurrents can enhance or interfere with colour  
 122 perception (e.g. [28,29]), and demonstrations of bidirectional-  
 123 ity in synaesthetic associations suggest that colour  
 124 concurrents can even affect the perception of the associated  
 125 trigger stimulus (e.g. [30]). However, consistent evidence for  
 126 a more generalized benefit of synaesthesia on multisensory

perception has not been forthcoming (see [31,32] for a discus-  
 sion). This is somewhat unexpected, given the evidence  
 for broad-range differences in patterns of cortical connectivity  
 and functional activation between synaesthetes and  
 non-synaesthetes ([33,34] but see [35]) including activation  
 differences in early visual regions of the brain [36]. Further-  
 more, given that synaesthesia is considered a 'mixing of the  
 senses', and involves multisensory processing of the inducing  
 stimulus [37], the specificity of the effect of the synaesthetic  
 experience on associated perceptual processes is surprising.  
 However, one possible reason for our lack of knowledge is  
 that the role of imagery in mediating cross-modal processes  
 has not been fully considered in investigations of synaesthe-  
 sia. Here, we conducted two experiments to assess whether  
 synaesthesia is associated with enhanced imagery abilities  
 and whether this has an impact on cross-modal perception.

## 2. Experiment 1: self-reported imagery in mirror- touch synaesthetes

MTS provides a candidate model of cross-modal synaesthesia  
 for investigating these effects: in MTS, the concurrent is  
 experienced in a sensory modality (touch) that differs from  
 that of the inducing stimulus (vision). An individual with  
 MTS automatically and consistently experiences a physical  
 sensation of touch on their own body when viewing touch  
 on another's body [38,39]. MTS has a reported prevalence  
 of approximately 1.6% [38] in the general population, and is  
 associated with more typical processes underpinning social  
 perception, including empathy [40]. The unique combination  
 of activation within mirror neuron systems in the brain,  
 which typically function to represent another person's actions  
 as one's own [41,42], as well as multisensory regions govern-  
 ing the sense of self, indicates that atypical self-processing and  
 self-other distinctions are present in MT synaesthetes [39].

We tested self-reported imagery in a group of MT  
 synaesthetes and compared their performance to two differ-  
 ent groups, a synaesthete group who did not have MTS  
 and a non-synaesthetic control group. Following an initial  
 screening phase, imagery in the tactile domain was tested  
 across two different tasks, somatosensory imagery (ratings  
 of sensitivity across the body) and tactile imagery (of object  
 properties) in all participants.

### (a) Methods

#### (i) Participants

Thirty participants volunteered to take part in this study and  
 were recruited from a large sample of individuals ( $N=366$ ,  
 described in [43]) who attended a public exhibition at the  
 Science Gallery Dublin. All participants were fluent English  
 speakers and 66% (20/30) reported English as their native  
 language. Based on our screening procedure (described  
 later), 10 individuals were identified as mirror-touch  
 synaesthetes (MT synaesthesia), 10 as having other forms of  
 synaesthesia (non-MT synaesthesia) and 10 as having no  
 synaesthete experiences (control). Both the non-MT  
 synaesthesia and control groups were age and sex matched  
 to the MT synaesthesia group, and selected via stratified  
 random sampling from larger samples of qualifying partici-  
 pants. Demographic details of all three groups of  
 participants are provided in table 1.

Q1

**Table 1.** Demographic information of the three groups (MTS, non-MT synaesthete and control) tested in Experiment 1.

	MT synaesthesia <i>N</i> = 10 (%)	non-MT synaesthesia <i>N</i> = 10 (%)	control <i>N</i> = 10 (%)
sex			
female	6 (60)	6 (60)	6 (60)
male	4 (40)	4 (40)	4 (40)
age			
13–17	3 (30)	3 (30)	3 (30)
18–35	4 (40)	4 (40)	4 (40)
36–55	2 (20)	2 (20)	2 (20)
56+	1 (10)	1 (10)	1 (10)
handedness			
right-handed	7 (70)	9 (90)	8 (80)
left-handed/ambidexterous	1 (10)/2 (20)	1 (10)	1 (10)/1 (10)
nationality			
European	6 (60)	9 (90)	8 (80)
North American	3 (30)	0	2 (20)
Asian	1 (10)	1 (10)	0
diagnosis of neurological disorder			
no	9 (90)	10 (100)	10 (100)
MT synaesthesia ratings (1–7)			
mean frequency of MTS	5.7	3	2.4
mean strength of MTS to others			
familiar	5.71	3.8	2.6
unfamiliar	4.29	2.1	1.8
additional types of synaesthesia (frequency of 'yes' response)			
none	1 (10)	0	10 (100)
grapheme–colour	2 (20)	2 (20)	0
music–colour	3 (30)	4 (40)	0
taste–colour	1 (10)	2 (20)	0
other (not listed)	3 (30)	2 (20)	0
ratings to mirror-touch videos (1–7)			
mean sensitivity (s.d.)	5.30 (1.26)	2.54 (1.57)	2.45 (1.72)
range	4.96–5.79	1–4.92	1–4.54
body location mean sensitivity (s.d.)			
back of neck	5.85 (1.27)	2.55 (1.55)	2.78 (2.04)
ear	5.12 (1.29)	2.77 (1.70)	2.58 (1.73)
hand (ego)	4.95 (1.19)	2.55 (1.64)	1.85 (1.09)
hand (allo)	5.63 (1.19)	2.77 (1.73)	2.52 (1.82)
mean sensitivity to stimulus type (s.d.)			
real hand	5.46 (1.22)	2.71 (1.70)	2.35 (1.68)
fake hand	5.04 (1.31)	2.51 (1.53)	2.45 (1.68)
paintbrush	5.40 (1.23)	2.40 (1.48)	2.55 (1.80)

**(ii) Materials and apparatus**

A general questionnaire was designed to determine basic demographic details of the participants (e.g. sex, age, nationality) as shown in table 1. The main study comprise three different tasks; a screening phase, somatosensory mental imagery (SMI) and tactile mental imagery (TMI) tasks. These two

tasks were conducted in a testing booth, measuring 2 m by 2 m, and surrounded by a black curtain that was closed during testing. The booth was positioned within a quiet section of the exhibition area of the Science Gallery. The entire experiment was presented on a computer (Dell Optiplex 790) with the monitor (1920 × 1080 pixel resolution with a

refresh rate of 60 Hz) resting on a table approximately 70 cm from the seated participant. All experimental tasks were programmed using Psychopy [44].

The screening phase involved both a questionnaire and rating study. The questionnaire included two questions relating to MTS specifically, and one question on other forms of synaesthesia. The MTS questions related to how frequently the participant experiences a sensation of touch on their own body when viewing touch on another person's body in everyday life (from 1 *never* to 7 *all the time*) and the strength of tactile mirroring when viewing touch on the body of a personally familiar or unfamiliar individual (from 1 *very weak* to 7 *very strong sensation*). Finally, participants were provided with a brief description of synaesthesia (*Synaesthesia is described as a curious mingling of the senses. For example, some people see colour or experience taste when they hear sounds*) and were asked to indicate if they always experience colour to any of the following stimuli (i) letters and/or numbers (i.e. indicating grapheme-colour synaesthesia), (ii) music (auditory-colour synaesthesia), (iii) while eating (gustatory-colour synaesthesia), or if they experienced (iv) another form of synaesthesia not listed. Participants were also asked to confirm (v) if they had synaesthesia. Participants provided 'yes' or 'no' confirmatory responses only to the general questions on synaesthesia.

For the rating study of the screening phase, the stimulus set comprise short video clips of one of two different models, a male and a female, being stroked on either their right hand (dorsal), their right ear or back of the neck. The hand was presented twice, in separate video clips, from an allocentric and egocentric perspective. Each body location was shown stroked three times per video clip, once each with a different stimulus inducer of either another person's real hand (opposite sex to the model), a fake hand or a paintbrush. The actor's hand controlling the fake hand or paintbrush was not visible during the video. The direction and velocity of the stroking were matched across videos to an approximate stroking speed of 8 cm s<sup>-1</sup>. A total of 24 video clips (two models, four body sites and three stimulus inducers) comprised the set of stimuli for the screening phase. Each video clip lasted no longer than 6 s and participants were presented with instructions to provide a rating response after viewing each clip.

For the SMI test, the stimulus set comprise schematic illustrations of individual body parts with specific anatomical regions highlighted in colour (for examples of these stimuli, see electronic supplementary material). The total number of 21 stimuli consisted of the following body sites: hallux, ball of the foot, foot underside, dorsal foot, the shin, thigh, abdomen, chest, neck, mouth, nose, cheek, ears, forehead, upper arm, forearm, dorsal hand, palm of the hand, dorsal finger, finger underside and the fingertips. Each stimulus was displayed for 2 s and participants were then presented with instructions to provide a rating response.

The stimuli for the TMI test consisted of the names of five objects; bubble wrap, ice cube, sandpaper, velvet and wet sponge. Each object name was presented at the top of the screen along with a specific instruction to rate the object on each of its properties (for an example of the stimulus display, see electronic supplementary material). These properties included surface force (e.g. required to pop the bubble wrap, to feel the ice or sandpaper etc.), resistance (hard or soft), texture (rough or smooth) and object weight (light or

heavy). Each name stimulus in this task was presented until the participant provided a rating response.

### (iii) Design and procedure

Before entering the testing booth, participants were first invited to take part in the study by providing informed, written consent and basic demographic information at a sign-in station. The parent or guardian of any individual under the age of 18 was required to complete the consent form before the individual could participate in the study. The demographic details of all participants are provided in table 1. Each participant was then escorted to the testing booth, where they were presented with the series of three rating tasks; the screening test for MTS, TMI and SMI tasks. Trials were randomly presented across participants for each of the imagery tasks. A practice trial was first provided that required the participant to rate the taste of a lemon (from 1 *very sweet* to 7 *very bitter*) using the same rating scale as in the other tasks. The screening test trials appeared immediately after this initial practice trial. Each video clip in a trial was preceded by a short, 3 s prompt which asked *How much do you feel this touch on your own body?* to which participants responded on a rating scale from 1 (*I felt no touch*) to 7 (*I felt the touch very strongly*).

For the TMI rating task, participants were instructed to imagine feeling various object materials (bubble wrap, ice cube, sandpaper, velvet, wet sponge) with their eyes open and to provide imagery ratings for each material (see electronic supplementary material for further details). Specifically, for each material, participants were required to indicate the vividness of their tactile imagery for all of the following object properties: force, resistance, texture and weight, using a Likert rating scale. Brief explanations were provided for these tactile properties, for example, participants were asked to imagine the force needed to pop the bubble wrap, imagine squeezing the wet sponge or imagine feeling the resistance (i.e. the hardness or softness) of a material. They were encouraged to provide ratings of their imagery ranging from 1 (*I cannot imagine this*) to 7 (*I can vividly imagine this*).

For the SMI rating task, participants were asked to refer to each schematic illustration of the body shown in each trial, and imagine being stimulated on their own corresponding body site by a light brushstroke. They then provided a rating of their sensitivity using a Likert scale ranging from 1 (*not sensitive at all*) to 7 (*extremely sensitive*).

To respond (in all rating tasks), the participant used a mouse to move a triangular marker along a scale until it reached the point on the scale corresponding to their rating. They then clicked the mouse to confirm their rating. The default position of the marker was always the midpoint of the scale and participants were encouraged to use the whole scale.

The final sequence of testing involved a series of questions which acted as a further screening test for synaesthesia, including two questions on MTS to which the participant provided a rating response, and five on other types of synaesthesia (or none), as described earlier. The entire experiment lasted no more than 10 min per participant.

### (iv) Data analysis

The analyses were performed using R (v. 3.5.0; [45]) on R studio [46]. The ANOVAs were conducted using the EZ

package [47] and type 3 sum-of-squares to test for significant main effects and interactions. Participants' mean Likert ratings to the screening, TMI and SMI tasks were entered into mixed-design ANOVAs for statistical analysis. Where appropriate, the Greenhouse–Geisser correction was applied to adjust the degrees of freedom of within-subject tests to correct for violations of the sphericity assumption and, in these cases, the adjusted  $p$ -value is reported. Separate post hoc tests were carried out for significant main effects or interactions (one-sample  $t$ -test for within-subject and Tukey HSD for between-subject comparisons). When multiple one-sample  $t$ -tests were performed, the Bonferroni correction was used to maintain a family-wise Type 1 error rate at 0.05 and the adjusted  $p$ -value is reported.

#### (v) 'MT' synaesthetes, 'non-MT' synaesthetes and 'control' group selection

The synaesthete participants were identified from their responses to the combined screening tests, that is, based on their self-reported answers to the series of questions on MTS and other types of synaesthesia, as well as their rating scores to the video clips designed to evoke MTS. We used two main, co-occurring inclusion criteria for MTS. First, we decided that, for each participant, the minimum cut-off rating in response to the mirror-touch videos was 5 (i.e. corresponding to a strong tactile sensation when viewing the videos) and the reported frequency of tactile mirroring experienced in everyday life should be higher than a rating of 4 (corresponding to 'a lot of the time'). Based on the overall mean rating scores in response to the videos clips alone, 12 candidates (3.3% of the total sample) qualified for consideration. Of these, three failed to meet our inclusion criterion of frequently experiencing mirror-touch (all three reported 'never' to whether they experienced a sensation of touch on their own body when viewing touch on other bodies in everyday life). We decided that an additional participant with a mean score of 4.96 in response to the video clips that was slightly lower than criterion, but who reported frequently experiencing MTS qualified for inclusion, yielding a total sample of 10 (MT synaesthesia group, table 1). This final sample of mirror-touch synaesthetes represents a prevalence rate of 2.7% from the larger sample, which is equivalent to, albeit slightly higher than, previous estimates of between 1.6 and 2.5% [37]. We also tested for the presence of co-occurring synaesthetic experiences in these and other participants. Nine participants from this MT synaesthesia group also reported experiencing another form of synaesthesia. All reported no history of physical impairment but one reported a history of neurological impairment (which we deemed as transient and unrelated to their synaesthesia).

The participants allocated to the MT synaesthesia group provided a mean rating of 5.70 ( $\pm 0.95$ ) on the frequency of tactile mirroring experienced in everyday life. Of these participants, seven provided mean ratings of 5.71 ( $\pm 0.95$ ) and 4.29 ( $\pm 2.06$ ) in response to familiar and unfamiliar individuals, respectively, which did not differ statistically [ $t_6 = 1.70$ ,  $p = 0.14$ ] (three MT synaesthete participants did not provide a response to this question). We found no evidence that the ratings to the different body sites, or stimulus types differed for this group (see 'Ratings to mirror-touch videos' in table 1).

Participants were allocated to the 'Non-MT synaesthesia' group or the 'control' group based on their responses to the


questionnaire (i.e. whether they reported experiencing synaesthesia or not), and on providing lower ratings to the video clips than the cut-offs mentioned earlier. Of the total number of synaesthetes identified through self-report ( $N = 165$ ), 10 were selected through stratified random sampling, with the constraints that they were age and sex matched to the MTS group, and had no history of neurological or physical impairments. The final sample included two grapheme–colour, four auditory–colour, two gustatory–colour and two reporting 'other' types of synaesthesia. Finally, 10 participants who self-reported experiencing no form of synaesthesia were selected through stratified random sampling and assigned to the control group. Again, these participants were matched to the MT synaesthesia group and reported no history of neurological or physical impairment. All ratings provided by each of these groups to the screening test (video clips) were significantly lower than those provided by the MTS group, as expected with our procedure ( $F_{2,27} = 20.4$ ,  $p < 0.001$ ,  $\eta_{\text{gen}}^2 = 0.5$ ). We also found a significant main effect of body location on the ratings ( $F_{2,54} = 4.96$ ,  $p = 0.015$ ,  $\eta_{\text{gen}}^2 = 0.02$ ), which was driven by higher ratings to the neck area than other body sites. No other differences or interactions were found. Further details of this analysis across groups and body location are provided in electronic supplementary material.

#### (b) Results


The mean ratings provided by each of the MT synaesthesia, non-MT synaesthesia and control groups to each of the tactile imagery (TMI) and somatosensory imagery (SMI) tasks are shown in figure 1. We first compared the groups' ratings to the TMI task. A mixed-model 3 (Group)  $\times$  4 (Object Property)  $\times$  5 (Material) ANOVA was performed on participants' mean ratings. A main effect of group was found ( $F_{2,27} = 4.86$ ,  $p < 0.01$ ,  $\eta_{\text{gen}}^2 = 0.15$ ). A post hoc Tukey HSD test revealed that the MT synaesthesia group provided significantly higher TMI ratings ( $M = 5.93 \pm 1.56$ ) than the non-MT synaesthesia ( $M = 4.72 \pm 1.85$ ;  $p < 0.001$ ) and control ( $M = 4.25 \pm 1.94$ ;  $p < 0.001$ ) groups, and the non-MT synaesthesia group provided higher ratings than the control group ( $p < 0.02$ ) (figure 1a).

There were also significant main effects of Material ( $F_{4,108} = 2.59$ ,  $p < 0.041$ ,  $\eta_{\text{gen}}^2 = 0.02$ ) and Object Property ( $F_{3,81} = 14.49$ ,  $p < 0.001$ ,  $\eta_{\text{gen}}^2 = 0.04$ ) which were qualified by a significant interaction between these factors ( $F_{12,324} = 6.54$ ,  $p < 0.001$ ,  $\eta_{\text{gen}}^2 = 0.05$ ) as shown in figure 2. Post hoc pairwise comparisons, with the Bonferroni correction, revealed differences between object properties to the sandpaper material only, with higher ratings to texture than force, lower ratings to weight than either texture or resistance and lower ratings to force than resistance (see electronic supplementary material, figure S1 for an illustration of these comparisons). Differences between material type were also found to be dependent on object properties but were mainly due to higher ratings on force and weight given to sandpaper than to the wet sponge or bubble wrap. The effect of group did not interact with any of the other factors.

The groups' mean ratings to the somatosensory imagery task are shown in figure 1b, and mean ratings provided by each group to each body site are provided in figure 3. To simplify the analysis, the 21 body parts were clustered into five main body regions as follows: head (mouth, nose,



**Figure 1.** The mean ratings provided to the (a) TMI and (b) SMI tasks by each of the control, non-MT synaesthesia (non-MT-Syn) and mirror-touch synaesthesia (MT-Syn) groups in Experiment 1. Individual participant mean ratings are also shown (as filled circles) within each group. Error bars represent 95% confidence intervals. (Online version in colour.)



**Figure 2.** Plot showing the interaction between ratings provided to materials and object properties found in Experiment 1. Error bars represent 95% confidence intervals. (Online version in colour.)


cheek, ears, forehead), upper body (abdomen, chest, neck, upper arm, forearm), lower body (shin, thigh), foot (hallux, dorsal foot, foot underside, ball of foot) and hand (dorsal hand, palm of the hand, dorsal finger, finger underside, fingertip). A mixed-model 3 (Group)  $\times$  5 (Body region) ANOVA was then conducted on participants' mean ratings to imagined sensitivity to a tactile stimulus on each viewed body site. A main effect of group was found ( $F_{2,27}=7.92$ ,  $p < 0.01$ ,  $\eta_{\text{gen}}^2 = 0.18$ ) and a post hoc Tukey HSD test confirmed significantly higher ratings by the MT synaesthesia group ( $M = 5.41 \pm 1.61$ ) than either the non-MT synaesthesia ( $M = 4.29 \pm 1.72$ ) or control ( $M = 4.66 \pm 1.48$ ) groups ( $ps < 0.001$ ), and lower ratings by the non-MT synaesthesia than control group ( $p = 0.047$ ). A main effect of body region was also found ( $F_{4,108} = 3.35$ ,  $p < 0.02$ ,  $\eta_{\text{gen}}^2 = 0.07$ ). Post hoc pairwise comparisons with the Bonferroni correction confirmed higher ratings to the head region ( $M = 5.16 \pm 0.87$ ) than the hand

( $M = 4.59 \pm 1.04$ ;  $p < 0.001$ ) and foot ( $M = 4.45 \pm 1.16$ ;  $p = 0.02$ ) regions. There was no evidence for an interaction between group and body region ( $p > 0.05$ ).

### (c) Discussion

Our results support the idea that synaesthesia is associated with enhanced imagery within the sensory domain of the synaesthetic experience [12], and provide evidence for cross-modal, visual to somatosensory imagery abilities. First, we found higher ratings of imagined touch in mirror-touch synaesthetes than in other synaesthetes or non-synaesthete controls when viewing regions of the body. This finding is not that surprising, given that mirrored somatosensory sensitivity when viewing tactile stimulation of another person is central to the reported experiences of mirror-touch synaesthetes [48], as well as reports of better tactile perception in this group [19]. Interestingly, the ratings of individuals with other reported forms of synaesthesia, so-called non-MT synaesthetes (mainly synaesthesia within visual and auditory modalities) were lower in the SMI test than those provided by the control group. This result is consistent with other reports of deactivated or negative neural responses in synaesthesia [34] and in imagery [49]. The difference between non-MT synaesthetes and controls was marginal, however, and future studies may help establish the veracity of this finding.

Our second finding, that MTS is associated with enhanced tactile imagery of object properties and materials that are not directly implicated in the synaesthetic experience itself, extends previous findings of enhanced imagery to within-modal stimuli beyond those experienced in synaesthesia [5,12]. Moreover, other non-MTS synaesthetes also reported enhanced imagery to these object stimuli relative to controls, suggesting a role for either multiple types of synaesthesia [12] or the involvement of vision in synaesthetic subtypes on enhanced tactile imagery. Indeed, we found it somewhat surprising that enhanced imagery for touch would extend to visual descriptions of distal objects and their properties in MTS in particular, given that such a task is more related to haptic perception than somatosensation *per se* [50]. However, touch can activate visual brain regions underpinning object



**Figure 3.** Participants mean ratings to the SMI task for each of the control, non-MT synaesthesia and MT synaesthesia groups (adapted from [48]). Groups are colour coded in different shades as follows: control (light), non-MT synaesthesia (intermediate) and MT synaesthesia (dark). The five main body regions are colour coded as follows: head (purple), upper body (yellow–orange), lower body (red), foot (blue) and hand (green). (Online version in colour.)

perception (i.e. the lateral occipital complex, see [51]); therefore, this finding may reveal the multisensory neural architecture supporting the representations of objects.

An important limitation to our study is that self-reported synaesthetes were not subjected to consistency testing which is often considered a hallmark of genuine synaesthesia [52]. We were constrained on the testing time available within the large public exhibition. Nevertheless, we are confident that our combined procedures helped us correctly identify individuals with MTS. Other types of synaesthetes were identified by questionnaire only, which we recognize as being less than ideal. Indeed, the prevalence of synaesthesia in our total sample was rather high (45% self-reported as synaesthetes) compared to previous estimates. This may have occurred if synaesthetes were particularly drawn to a public exhibition entitled ‘Fake’ which was advertised as also including tests of perception. However, other results assure us that these cases are also genuine. First, the prevalence rates from our sample are consistent with the literature in that more synaesthetes identified as having grapheme–colour synaesthesia than other forms [5,53] and MTS was a relatively rare form of synaesthesia [54]. Second, the reported main effects did not change when we repeated the stratified, random-sampling approach on the non-MT synaesthesia and control groups.

### 3. Experiment 2: cross-modal correspondences from the synaesthetic colours

The results of Experiment 1 suggest that synaesthesia is associated with imagery within the tactile domain that is triggered by a stimulus in another modality (vision). Unlike other cross-modal associations that are common to all [55], and influenced by associations found in the natural environment (e.g. [56]), it is assumed that synaesthetic experiences are resistant to learning. In other words, although the specific synaesthetic


associations may initially arise through learning (e.g. [57,58]), once the concurrent is established, the specific synaesthetic association with the inducer persists despite repeated evidence to the contrary from the environment (cf. changes due to ageing, [59]; or pharmacological intervention, [60]).

However, the assumption that synaesthetic experiences are encapsulated is not consistent with evidence that synaesthesia can affect perception, recall or even enhanced cross-modal imagery reported earlier. It remains possible, therefore, that the synaesthetic concurrent can, in itself, tap into other forms of information processing, although evidence for such influences beyond the synaesthetic association (e.g. letters and colours in grapheme colour synaesthesia) remains elusive. Nevertheless, such evidence might help provide insight into the mechanisms supporting the general enhancements associated with synaesthesia. Here, we investigated the extent to which synaesthetic experiences, or concurrents, are encapsulated using a method which exploits cross-modal correspondences. We were particularly interested in investigating whether synaesthesia influences multisensory or cross-modal processes, since these reflect more typical perceptual tasks in the natural environment, as well as neural processing in the brain [61]. Specifically, we tested whether cross-modal, tactile correspondences were also associated with synaesthetic experiences in the visual domain. To that end, we adapted a task described by Pinkerton & Humphrey [62] which was designed to test whether different colours are associated with differences in the tactile property of weight [63].

#### (a) Methods

##### (i) Participants

Sixteen grapheme–colour synaesthetes (12 female, mean age: 21.53 years old) and 22 age-matched controls (16 female, mean age: 22.1 years old) with normal or corrected-to-normal



**Figure 4.** (a) Example of a stimulus in the colour patch test used in Experiment 2. (b) Mean weight ratings provided across both groups of participants to colour patches. Error bars represent 95% confidence intervals. (Online version in colour.)

vision volunteered to take part in the study. Participants were recruited from the student population of Trinity College Dublin and were compensated with research credits for their time. All participants were naive to the purposes of the study and gave their informed, written consent prior to their inclusion.

All participants in the synaesthesia group reported seeing colours in response to letter and digit stimuli. Synaesthesia was verified using the Synaesthesia Test Battery [64] and a customized test of consistency matching over time (see *Procedure*). Participants with a score of less than 0.9 on the test battery and a test–retest consistency of 80% over a period of at least one week on the customized test were included in the synaesthesia group. Participants who failed to meet either of these criteria were excluded from the study. Control participants also performed both tests to rule out the possibility that they were subject to synaesthetic experiences.

### (ii) Stimuli

Stimuli consisted of circular colour patches and uppercase letters presented on a grey background ( $13 \text{ cd m}^{-2}$ ). Patch stimuli were masked by a Gaussian luminance profile with a fixed standard deviation of  $1.32^\circ$  and could take one of six isoluminant colours (red, blue, green, orange, yellow, white) with an average luminance of ( $13 \text{ cd m}^{-2}$ ). Letter stimuli were presented in Times New Roman font and were  $8^\circ$  of visual angle in height from a viewing distance of 57 cm. Fifteen-letter stimuli were used in total, all of which were white. Stimuli were displayed on a  $\gamma$ -corrected BenQ XL2410T LCD monitor at a resolution of  $1280 \times 1024$  pixels at a refresh rate of 120 Hz. The maximum luminances available for red, green and blue were 56, 178 and  $13 \text{ cd m}^{-2}$ , respectively. All stimuli were created and all data were collected using PsychoPy [44,65].

### (iii) Procedure

Participants performed a weight-matching task with colour patch and letter stimuli in separate blocks, the order of which was counter-balanced across participants. In both blocks, participants were presented with two stimuli located  $8^\circ$  either side of the centre of the screen and a rating scale located below the stimuli (figure 4a). Participants were instructed to imagine that the mouse cursor was an adjustable fulcrum and to position it at the point on the rating scale at which they judged the two stimuli to be balanced in weight.


There was no time limit on responses and once the participant was satisfied with their answer, they clicked an ‘Accept’ button. Prior to the experiment, participants completed a short practice block to familiarize the participant with the task.

Each participant completed a total of 200 weight-matching judgements (100 trials with colour patch and 100 trials with letter stimuli). In the colour patch block, the white stimulus acted as the standard stimulus and was present for all trials. The test stimulus was selected at random on each trial from the remaining colours, with each test colour being displayed 20 times. The positions of the standard and test stimuli were randomized on each trial, such that there was an equal probability that the standard stimulus would appear on the left or right of the screen. The procedure for the colour patch block was identical for the synaesthesia and control groups.

The letter stimuli were determined from a customized test administered to the synaesthesia group prior to the experiment in which participants provided a detailed description of the colour induced by each letter. The test was repeated two weeks later, to ensure retest consistency of 85% or above. To select appropriate letter stimuli for each participant, letter descriptions that included any of the colours used in colour patch block were initially identified as potential stimuli for the letter block and, where possible, letters were retained when the induced colour was also described as ‘strong’ or ‘saturated’. To select a standard stimulus for each participant, preference was given to letters that either did not induce a colour or induced the colour white. This letter selection process led to a unique set of letters for each synaesthete, with 15 distinct letters used across all 10 participants. For each participant in the control group, five test letters were selected at random from the 10 most frequently selected letters in the synaesthesia group, while the standard letter was always the letter ‘O’. All other aspects of the experimental procedure were identical to the colour patch block.

### (iv) Data analysis

Individual weight judgements were converted to a numeric value between 5 and  $-5$ , where positive/negative values indicated that the test stimulus was judged to be heavier/lighter than the standard. Each test stimulus was presented 20 times in a block and the mean of these weight judgements was taken as the weight rating for that stimulus. These values were subsequently group-averaged for statistical analysis,



**Figure 5.** 'Weight' responses in synaesthete group to letters and colour patches. Error bars represent 95% confidence intervals.

where within- and between-group comparisons were made using different types of ANOVA. Where appropriate, the Greenhouse–Geisser correction was applied to adjust the degrees of freedom of within-subject tests to correct for violations of the sphericity assumption and, in these cases, the adjusted  $p$ -value is reported. When multiple one-sample  $t$ -tests were performed, the Bonferroni correction was used to maintain a family-wise Type 1 error rate at 0.05 and the adjusted  $p$ -value is reported. All statistical analyses were performed in GraphPad Prism 6.0, with type 3 sum-of-squares used to test for significant main effects and interactions.

## (b) Results

The group-averaged results for the synaesthesia group and the age-matched controls to the colour patches are shown in figure 4*b*. For both groups, all colours tested were judged to be heavier than the standard white stimulus, as indicated by the positive weight ratings for every test colour. Furthermore, the magnitude of the weight rating varied for different colours, with a similar ranking of weights observed across both groups. A one-way repeated-measures ANOVA confirmed that the magnitude of weight judgements varied for different colours in both the control group ( $F_{3,74} = 38.18$ ,  $p_{\text{adj}} < 0.001$ ,  $\eta^2 = 0.45$ ) and the synaesthesia group ( $F_{3,49} = 35.76$ ,  $p_{\text{adj}} < 0.001$ ,  $\eta^2 = 0.58$ ), while one-sample  $t$ -tests against a weight rating of 0 determined that all colour patches were significantly heavier than the white stimulus (control group: ( $t_{521} > 6.12$ ,  $p_{\text{adj}} < 0.001$ ), synaesthesia group: ( $t_{15} > 6.97$ ,  $p_{\text{adj}} < 0.001$ )). Finally, a mixed-model 2 (group)  $\times$  5 (colour) ANOVA revealed that there was a significant main effect of group on weight ratings ( $F_{1,36} =$ ,  $p < 0.001$ ,  $\eta^2 = 0.17$ ), with synaesthetes ranking the colours as heavier overall.

We then examined whether the colours experienced by grapheme–colour synaesthetes in response to letters also have an apparent weight. To facilitate comparison with the colour patch data, we averaged across these different letter–colour combinations, such that the data in figure 5 are expressed in terms of the weight responses to the colour patches (Colour) and to the colour induced by the letter stimuli (hereafter referred to as 'letter-induced colours'). As with the colour patch data, all of the letter-induced colours were rated as heavier than the standard stimulus, as indicated by the positive weight rating values shown in figure 5. Furthermore, there was a striking resemblance in the ranking and the relative magnitudes of weight ratings for both

datasets, with red and blue again being ranked as the heaviest and yellow as the lightest. A one-way repeated-measures ANOVA confirmed that the magnitude of weight ratings varied for the different induced colours ( $F_{3,43} = 16.54$ ,  $p_{\text{adj}} < 0.001$ ,  $\eta^2 = 0.3$ ), although subsequent one-sample  $t$ -tests revealed that only the red, blue and green conditions were significantly different from a weight rating of 0 (red:  $t_{15} = 6.17$ ,  $p_{\text{adj}} < 0.001$ , Cohen's  $d = 1.55$ ; blue:  $t_{15} = 9.38$ ,  $p_{\text{adj}} < 0.001$ , Cohen's  $d = 2.34$ ; green:  $t_{15} = 2.97$ ,  $p_{\text{adj}} < 0.05$ , Cohen's  $d = 0.74$ ). The weight rating responses to the colour patch stimuli were significantly higher than those to the letter stimuli ( $F_{1,30} = 4.887$ ,  $p = 0.035$ ,  $\eta^2 = 0.076$ ). We also conducted a correlation analyses which revealed a strong positive correlation between the weight ratings for the two types of stimuli in the synaesthete group ( $R^2 = 0.372$ ,  $p < 0.001$ ), while the slope of the best fitting regression line to the data was significantly different from 0 ( $a = 0.361$ ,  $p < 0.001$ ). Moreover, fitting individual regression lines to each participant's data confirmed that there was a positive slope for all 16 datasets (ranging between 0.22 and 1.67), suggesting that this relationship was robust to the level of individual participants. Crucially, we found no consistent pattern in the weight ratings assigned to different letters by the control group and a one-sample  $t$ -test confirmed that overall these ratings were not significantly different from a weight rating of 0 ( $t_{109} = 0.9466$ ,  $p = 0.35$ , Cohen's  $d = 0.09$ ).

## (c) Discussion

First, our results based on real colour patches replicated those previously reported [62]: red and blue were consistently perceived as the heaviest colours, and yellow as the lightest. Moreover, our findings provide an important extension: here, we found that induced, synaesthetic colours are sufficient to evoke cross-modal correspondences of weight, typically considered a tactile property of objects, and do so in a manner that is akin to the correspondences experienced with veridical colours. For grapheme–colour synaesthetes, letters that induced the colour red, or blue, were perceived as 'heaviest' while letters that induced the colour yellow were perceived as lighter. This finding challenges the idea that the effect of the synaesthetic concurrent on perception is specific to its domain (e.g. colour, 18; or somatosensation, 19), or that of the inducer [30]. By contrast, our results provide evidence for the perceptual reality of the synaesthetic concurrent, particularly its role in cross-modal interactions that are commonly experienced.

## 4. General discussion

Our results add to growing evidence that synaesthesia is linked to more vivid imagery, and that imagery abilities generalize to stimuli not directly involved in the synaesthetic experience [5,12]. Furthermore, we provide evidence that MTS is associated with heightened cross-modal imagery: visual stimuli evoke proximal tactile sensations on the body as well as enhanced tactile imagery of distal, external objects. A distinction between passive (somatosensation) and active (haptics) touch has previously been reported within the tactile system [66]; therefore, these results are interesting in that they show generalization of imagery abilities beyond the limits of the synaesthetic interactions themselves as well as the functional organization of the somatosensory system.

We also report a novel finding that a visual synaesthetic concurrent can mediate cross-modal interactions. The results of our second experiment show that a tactile feature, weight, can be readily associated with an induced colour in a manner that is similar to viewing the real colour. Thus, correspondences were generated in a modality beyond vision that is not typically associated with grapheme–colour synaesthesia. Furthermore, the nature of the cross-modal correspondences were not unique to the grapheme–colour synaesthetes. Taken together, our findings challenge the idea that the synaesthetic concurrent is entirely encapsulated. Specifically, synaesthesia is associated with a general ability to imagine other stimuli within the same modality, as well as influencing perceptual associations in other modalities. It is unclear, therefore, why the synaesthetic concurrent appears to resist other cognitive influences and, despite the evidence that it is associated with individual differences in imagery and perception, appears to coexist with these other experiences without the representation of the synaesthetic concurrent being affected.

Although our study does not directly address the issue of causality, there may be more to the ‘suspicious coincidence’ that enhanced imagery is found in synaesthetes. For example, it may be that both synaesthesia [5,67] and vivid imagery [16,68] arise from higher baseline excitability within sensory cortices than in the general population. The cortical responses in S1 and SII in mirror-touch synaesthetes are particularly intriguing, first because they arise in response to visual depictions of tactile stimulation on another’s body and second, the responses are specific to viewing real bodies being touched as opposed to objects or fake bodies [14]. Moreover, hyper-excitability of S1, induced by neuro-stimulation, produces mirror-touch phenomena [3,4], and (pre-) viewing another person’s body region modifies tactile sensitivity on one’s own body [69], in non-synaesthetic populations. These findings suggest that similar cross-modal connections are present in all, synaesthete and non-synaesthete alike.

Our results also raise the question of what is the nature of the representations at the interface between imagery, synaesthesia and perception. Clearly, there are important phenomenological differences between synaesthesia and imagery, as well as the fact that imagery is more widely experienced. Previous studies suggest that any differences found between perception, imagery and synaesthesia may

reflect different stages of information processing along the sensory pathways. For example, Arnold *et al.* [70] reported that the variability in responses to synaesthetic colour matching was similar to colour matching from memory, but larger than to real colour matching, suggesting that synaesthetic colours are akin to colour memory than perception and that synaesthetic colours likely arise at a higher level of analysis in the brain [71]. Indeed, these psychophysical findings are consistent with neuroimaging reports demonstrating activation in higher areas of the brain to synaesthetic than imagined colours [72], suggesting more top-down influences in internally generated visual experiences [32,73]. It may be these top-down influences, in particular, that allow for the synaesthetic representation to persist despite competing information from the environment (see [74]).

In sum, although synaesthetic experiences are automatically generated and are consistent over time, our findings challenge the idea that induced synaesthetic experiences are fully cognitively encapsulated. Rather, synaesthesia can influence other cognitive processes including cross-modal perception. These findings have important implications for our understanding of the modular basis of synaesthesia.

**Ethics.** All procedures were approved by the School of Psychology Research Ethics Committee at Trinity College Dublin. Accordingly, participants gave informed, written consent prior to taking part in any of the studies described above.

**Data accessibility.** The datasets supporting this article will be uploaded as part of the electronic supplementary material.

**Authors’ contributions.** A.O.D., S.M.C. and F.N.N. contributed to the design of Experiment 1, A.O.D. collected the data and A.O.D., S.M.C. and F.N.N. analysed and interpreted the data. D.P.M.G. and F.N.N. contributed to the design of Experiment 2, and D.P.M.G. analysed and interpreted the data. All authors contributed towards drafting and revising the article. All approved the final version for publication.

**Competing interests.** We have no competing interests.

**Funding.** The research contained in this article was funded from a variety of sources including a TCD post-graduate scholarship (A.O.D.), EU-Commission, grant no. 732391 (S.M.C., F.N.N.) and a Government of Ireland Fellowship from the Irish Research Council (D.P.M.G.).

**Acknowledgements.** The data described here for Experiment 2 were collected during an exhibition (entitled ‘Fake’) held at Science Gallery Dublin in 2018. We thank Patrick McNestry for help in collecting the data for Experiment 2.

## References

- Galton F. 1907 *Inquiries into human faculty and its development*. London, UK: JM Dent & Sons.
- Ramachandran VS, Hubbard EM. 2001 Synaesthesia—a window into perception, thought and language. *J Consciousness Stud.* **8**, 3–34.
- Craver-Lemley C, Reeves A. 2013 Is synesthesia a form of mental imagery? In *Multisensory imagery* (eds S Lacey, R Lawson), pp. 185–206. New York, NY: Springer.
- Price MC. 2009 Spatial forms and mental imagery. *Cortex* **45**, 1229–1245. (doi:10.1016/j.cortex.2009.06.013)
- Barnett KJ, Newell FN. 2008 Synaesthesia is associated with enhanced, self-rated visual imagery. *Conscious Cogn.* **17**, 1032–1039. (doi:10.1016/j.concog.2007.05.011)
- Spiller MJ, Jansari AS. 2008 Mental imagery and synaesthesia: is synaesthesia from internally-generated stimuli possible? *Cognition* **109**, 143–151. (doi:10.1016/j.cognition.2008.08.007)
- Simner J, Mayo N, Spiller MJ. 2009 A foundation for savantism? Visuo-spatial synaesthetes present with cognitive benefits. *Cortex* **45**, 1246–1260. (doi:10.1016/j.cortex.2009.07.007)
- Brang D, Miller LE, McQuire M, Ramachandran VS, Coulson S. 2013 Enhanced mental rotation ability in time-space synesthesia. *Cogn. Process* **14**, 429–434. (doi:10.1007/s10339-013-0561-5)
- Havlik AM, Carmichael DA, Simner J. 2015 Do sequence-space synaesthetes have better spatial imagery skills? Yes, but there are individual differences. *Cogn. Process* **16**, 245–253. (doi:10.1007/s10339-015-0657-1)
- Mann H, Korzenko J, Carriere JSA, Dixon MJ. 2009 Time–space synaesthesia—a cognitive advantage? *Conscious Cogn.* **18**, 619–627. (doi:10.1016/j.concog.2009.06.005)
- Rizza A, Price MC. 2012 Do sequence-space synaesthetes have better spatial imagery skills? Maybe not. *Cogn. Process* **13**, 299–303. (doi:10.1007/s10339-012-0459-7)
- Spiller MJ, Jonas CN, Simner J, Jansari A. 2015 Beyond visual imagery: how modality-specific is

- 631 enhanced mental imagery in synesthesia? *Conscious*  
 632 *Cogn.* **31**, 73–85. (doi:10.1016/j.concog.2014.10.  
 633 010)
- 634 13. Ishai A, Sagi D. 1995 Common mechanisms of visual  
 635 imagery and perception. *Science* **268**, 1772–1774.  
 636 (doi:10.1126/science.7792605)
- 637 14. Ishai A, Sagi D. 1997 Visual imagery facilitates  
 638 visual perception: psychophysical evidence. *J. Cogn.*  
 639 *Neurosci.* **9**, 476–489. (doi:10.1162/jocn.1997.9.4.  
 640 476)
- 641 15. Marks DF. 1973 Visual imagery differences in the  
 642 recall of pictures. *Br. J. Psychol.* **64**, 17–24. (doi:10.  
 643 1111/j.2044-8295.1973.tb01322.x)
- 644 16. Kosslyn SM, Ganis G, Thompson WL. 2001 Neural  
 645 foundations of imagery. *Nat. Rev. Neurosci.* **2**, 635.  
 646 (doi:10.1038/35090055)
- 647 17. Ishai A, Ungerleider LG, Haxby JV. 2000 Distributed  
 648 neural systems for the generation of visual images.  
 649 *Neuron* **28**, 979–990. (doi:10.1016/S0896-  
 650 6273(00)00168-9)
- 651 18. Banissy MJ, Tester V, Muggleton NG, Janik AB,  
 652 Davenport A, Franklin A, Walsh V, Ward J. 2013  
 653 Synesthesia for color is linked to improved color  
 654 perception but reduced motion perception. *Psychol.*  
 655 *Sci.* **24**, 2390–2397. (doi:10.1177/0956797613492424)
- 656 19. Banissy MJ, Walsh V, Ward J. 2009 Enhanced  
 657 sensory perception in synaesthesia. *Exp. Brain Res.*  
 658 **196**, 565–571. (doi:10.1007/s00221-009-1888-0)
- 659 20. Alvarez BD, Robertson LC. 2013 The interaction of  
 660 synesthetic and print color and its relation to visual  
 661 imagery. *Atten. Percept. Psychophys.* **75**,  
 662 1737–1747. (doi:10.3758/s13414-013-0520-3)
- 663 21. Chiou R, Rich AN, Rogers S, Pearson J. 2018  
 664 Exploring the functional nature of synaesthetic  
 665 colour: dissociations from colour perception and  
 666 imagery. *Cognition* **177**, 107–121. (doi:10.1016/j.  
 667 cognition.2018.03.022)
- 668 22. Dixon M, Smilek D, Cudahy C, Merikle PM. 2000  
 669 Five plus two equals yellow. *Nature* **406**, 365.  
 670 (doi:10.1038/35019148).
- 671 23. Nikolić D, Jürgens UM, Rothen N, Meier B, Mroczko  
 672 A. 2011 Swimming-style synesthesia. *Cortex* **47**,  
 673 874–879. (doi:10.1016/j.cortex.2011.02.008)
- 674 24. Lacey S, Lawson R. (Eds.). 2013 *Multisensory*  
 675 *imagery*. Berlin, Germany: Springer Science &  
 676 Business Media.
- 677 25. Nanay B. 2018 Multimodal mental imagery. *Cortex*  
 678 **105**, 125–134. (doi:10.1016/j.cortex.2017.07.006)
- 679 26. Berger CC, Ehrsson HH. 2013 Mental imagery  
 680 changes multisensory perception. *Curr. Biol.* **23**,  
 681 1367–1372. (doi:10.1016/j.cub.2013.06.012)
- 682 27. Lacey S, Tal N, Amedi A, Sathian K. 2009 A putative  
 683 model of multisensory object representation. *Brain*  
 684 *Topogr.* **21**, 269–274. (doi:10.1007/s10548-009-  
 685 0087-4).
- 686 28. Smilek D, Dixon MJ, Cudahy C, Merikle PM. 2001  
 687 Synaesthetic photisms influence visual perception.  
 688 *J. Cogn. Neurosci.* **13**, 930–936. (doi:10.1162/  
 689 089892901753165845)
- 690 29. Palmeri TJ, Blake R, Marois R, Flanery MA, Whetsell  
 691 Jr W. 2002 The perceptual reality of synesthetic  
 692 colors. *Proc. Natl Acad. Sci. USA* **99**, 4127–4131.  
 693 (doi:10.1073/pnas.022049399)
30. Kadosh RC, Sagiv N, Linden DE, Robertson LC,  
 Elinger G, Henik A. 2005 When blue is larger than  
 red: colors influence numerical cognition in  
 synesthesia. *J. Cogn. Neurosci.* **17**, 1766–1773.  
 (doi:10.1162/089892905774589181)
31. Whittingham KM, McDonald JS, Clifford CW. 2014  
 Synesthetes show normal sound-induced flash  
 fusion and fusion illusions. *Vision Res.* **105**, 1–9.  
 (doi:10.1016/j.visres.2014.08.010)
32. Newell FN, Mitchell KJ. 2016 Multisensory  
 integration and cross-modal learning in  
 synaesthesia: a unifying model. *Neuropsychologia*  
**88**, 140–150. (doi:10.1016/j.neuropsychologia.2015.  
 07.026)
33. Rouw R, Scholte HS. 2007 Increased structural  
 connectivity in grapheme-color synesthesia. *Nat.*  
*Neurosci.* **10**, 792. (doi:10.1038/nn1906)
34. O'Hanlon E, Newell FN, Mitchell K. 2013 Combined  
 structural and functional imaging reveals cortical  
 deactivations in grapheme-color synaesthesia. *Front.*  
*Psychol.* **4**, 755. (doi:10.3389/fpsyg.2013.00755)
35. Hupé J-M, Bordier C, Dojat M. 2012 The neural  
 bases of grapheme-color synesthesia are not  
 localized in real color-sensitive areas. *Cereb. Cortex*  
**22**, 1622–1633. (doi:10.1093/cercor/bhr236)
36. Barnett KJ, Foxe JJ, Molholm S, Kelly SP, Shalgi S,  
 Mitchell KJ, Newell FN. 2008 Differences in early  
 sensory-perceptual processing in synesthesia: a  
 visual evoked potential study. *Neuroimage* **43**,  
 605–613. (doi:10.1016/j.neuroimage.2008.07.028)
37. Bargary G, Barnett KJ, Mitchell KJ, Newell FN. 2009  
 Colored-speech synaesthesia is triggered by  
 multisensory, not unisensory, perception. *Psychol.*  
*Sci.* **20**, 529–533. (doi:10.1111/j.1467-9280.2009.  
 02338.x)
38. Banissy MJ, Kadosh RC, Maus GW, Walsh V, Ward J.  
 2009 Prevalence, characteristics and a  
 neurocognitive model of mirror-touch synaesthesia.  
*Exp. Brain Res.* **198**, 261–272. (doi:10.1007/s00221-  
 009-1810-9)
39. Banissy MJ, Ward J. 2013 Mechanisms of self-other  
 representations and vicarious experiences of touch  
 in mirror-touch synesthesia. *Front. Hum. Neurosci.* **7**,  
 112. (doi:10.3389/fnhum.2013.00112)
40. Banissy MJ, Ward J. 2007 Mirror-touch synesthesia  
 is linked with empathy. *Nat. Neurosci.* **10**, 815.  
 (doi:10.1038/nn1926)
41. Rizzolatti G, Fogassi L, Gallese V. 2001  
 Neurophysiological mechanisms underlying the  
 understanding and imitation of action. *Nat. Rev.*  
*Neurosci.* **2**, 661. (doi:10.1038/35090060)
42. Gallese V. 2005 'Being like me': self-other identity,  
 mirror neurons and empathy. In *Perspectives on*  
*imitation: from cognitive neuroscience to social*  
*science* (eds SL Hurley, N Chater), pp. 101–118. MA:  
 MIT Press.
43. Cooney S, O'Dowd A, Newell F. 2019 Individual  
 differences in mental imagery abilities in the tactile  
 domain. Manuscript in preparation.
44. Peirce JW, MacAskill MR. 2018 *Building experiments*  
*in PsychoPy*. London, UK: Sage.
45. R Core Team. 2017 *R: a language and environment*  
*for statistical computing*. Vienna, Austria: R  
 Foundation for Statistical Computing. See <https://www.R-project.org/>.
46. RStudio Team. 2015 *RStudio: integrated*  
*development for R*. Boston, MA: RStudio, Inc. See  
<http://www.rstudio.com/>.
47. Lawrence MA. 2016 ez: easy analysis and  
 visualization of factorial experiments. R package  
 version 4.4-0. See [https://CRAN.R-project.org/  
 package=ez](https://CRAN.R-project.org/package=ez).
48. Weinstein S. 1968 Intensive and extensive aspects of  
 tactile sensitivity as a function of body part, sex and  
 laterality. In *The skin senses* (ed. DR Kenshalo).  
 Springfield, IL: Charles C Thomas.
49. Amedi A, Malach R, Pascual-Leone A. 2005 Negative  
 BOLD differentiates visual imagery and perception.  
*Neuron* **48**, 859–872. (doi:10.1016/j.neuron.2005.  
 10.032)
50. Lederman SJ, Klatzky RL. 2009 Haptic perception: a  
 tutorial. *Atten. Percept. Psychophys.* **71**, 1439–1459.  
 (doi:10.3758/APP.71.7.1439)
51. Amedi A, Malach R, Hendler T, Peled S, Zohary E.  
 2001 Visuo-haptic object-related activation in the  
 ventral visual pathway. *Nat. Neurosci.* **4**, 324.  
 (doi:10.1038/85201)
52. Cytowic RE. 1997 Synaesthesia: phenomenology and  
 neuropsychology—a review of current knowledge.  
 In *Synaesthesia: classic and contemporary readings*  
 (eds S Baron-Cohen, JE Harrison), pp. 17–39.  
 Malden, MA: Blackwell Publishing (Reprinted from  
 'Psyche', 1996).
53. Day S. 2001 Types of synesthesia. Web site on  
 synaesthesia maintained by Sean Day. See [http://  
 www.users.muohio.edu/daysa/types.htm](http://www.users.muohio.edu/daysa/types.htm).
54. Banissy MJ. 2013 Synaesthesia, mirror neurons and  
 mirror-touch. In *The Oxford handbook of*  
*synaesthesia* (eds E Hubbard, J Simner), pp.  
 584–605. Oxford, UK: Oxford University Press.
55. Spence C. 2011 Crossmodal correspondences: a  
 tutorial review. *Atten. Percept. Psychophys.* **73**,  
 971–995. (doi:10.3758/s13414-010-0073-7)
56. Parise CV, Knorre K, Ernst MO. 2014 Natural auditory  
 scene statistics shapes human spatial hearing. *Proc.*  
*Natl Acad. Sci. USA* **111**, 6104–6108. (doi:10.1073/  
 pnas.1322705111)
57. Watson MR, Akins K, Spiker C, Crawford L, Enns JT.  
 2014 Synesthesia and learning: a critical review and  
 novel theory. *Front. Neurosci.* **8**, 98. (doi:10.3389/  
 fnhum.2014.00098)
58. Witthoft N, Winawer J. 2013 Learning, memory,  
 and synesthesia. *Psychol. Sci.* **24**, 258–265. (doi:10.  
 1177/0956797612452573)
59. Simner J, Ipser A, Smees R, Alvarez J. 2017 Does  
 synaesthesia age? Changes in the quality and  
 consistency of synaesthetic associations.  
*Neuropsychologia* **106**, 407–416. (doi:10.1016/j.  
 neuropsychologia.2017.09.013)
60. Luke D, Terhune DB. 2013 The induction of  
 synaesthesia with chemical agents: a systematic  
 review. *Front. Psychol.* **4**, 753. (doi:10.3389/fpsyg.  
 2013.00753)
61. Ghazanfar AA, Schroeder CE. 2006 Is neocortex  
 essentially multisensory? *Trends Cogn. Sci.* **10**,  
 278–285. (doi:10.1016/j.tics.2006.04.008)

- 694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756
62. Pinkerton E, Humphrey NK. 1974 The apparent heaviness of colours. *Nature* **250**, 164. (doi:10.1038/250164a0)
63. Bullough E. 1907 On the apparent heaviness of colours: a contribution to the aesthetics of colour. *Br. J. Psychol.* **2**, 111–152. (doi:10.1111/j.2044-8295.1907.tb00236.x)
64. Eagleman DM, Kagan AD, Nelson SS, Sagaram D, Sarma AK. 2007 A standardized test battery for the study of synaesthesia. *J. Neurosci. Methods* **159**, 139–145. (doi:10.1016/j.jneumeth.2006.07.012)
65. Peirce, J. W. 2009 Generating stimuli for neuroscience using PsychoPy. *Front. Neuroinform.* **2**, 10. (doi:10.3389/neuro.11.010.2008)
66. Simões-Franklin C, Whitaker TA, Newell FN. 2011 Active and passive touch differentially activate somatosensory cortex in texture perception. *Hum. Brain Mapp.* **32**, 1067–1080. (doi:10.1002/hbm.21091)
67. Terhune DB, Tai S, Cowey A, Popescu T, Kadosh RC. 2011 Enhanced cortical excitability in grapheme-color synesthesia and its modulation. *Curr. Biol.* **21**, 2006–2009. (doi:10.1016/j.cub.2011.10.032)
68. Fallgatter AJ, Mueller TJ, Strik WK. 1997 Neurophysiological correlates of mental imagery in different sensory modalities. *Int. J. Psychophysiol.* **25**, 145–153. (doi:10.1016/S0167-8760(96)00738)
69. Kennett S, Taylor-Clarke M, Haggard P. 2001 Noninformative vision improves the spatial resolution of touch in humans. *Curr. Biol.* **11**, 1188–1191. (doi:10.1016/S0960-9822(01)00327-X)
70. Arnold DH, Wegener SV, Brown F, Mattingley JB. 2012 Precision of synesthetic color matching resembles that for recollected colors rather than physical colors. *J. Exp. Psych.: Hum. Percept. Perform.* **38**, 1078. (doi:10.1037/a0028129)
71. Erskine H, Mattingley JB, Arnold DH. 2013 Synaesthesia and colour constancy. *Cortex* **49**, 1082–1088. (doi:10.1016/j.cortex.2012.03.002)
72. Rich AN, Williams MA, Puce A, Syngeiotis A, Howard MA, McGlone F, Mattingley JB. 2006 Neural correlates of imagined and synaesthetic colours. *Neuropsychologia* **44**, 2918–2925. (doi:10.1016/j.neuropsychologia.2006.06.024)
73. Mechelli A, Price CJ, Friston KJ, Ishai A. 2004 Where bottom-up meets top-down: neuronal interactions during perception and imagery. *Cereb. Cortex* **14**, 1256–1265. (doi:10.1093/cercor/bhh087)
74. Grossenbacher PG, Lovelace CT. 2001 Mechanisms of synesthesia: cognitive and physiological constraints. *Trends Cogn. Sci.* **5**, 36–41. (doi:10.1016/S1364-6613(00)01571-0)