

Negative Phototaxis Behaviour of Organic Droplets in Channels

Larisa Florea¹, Klaudia Wagner², Pawel Wagner², Gordon W. Wallace²,
David L. Officer², Fernando Benito-Lopez^{1,3} and Dermot Diamond¹



¹The INSIGHT Centre for Data Analytics; National Centre for Sensor Research; School of Chemical Sciences, Dublin City University, Dublin 9, Ireland, ²ARC Centre of Excellence for Electromaterials Science and Intelligent Polymer Research, Institute University of Wollongong, Wollongong, Australia, ³CIC microGUNE, Arrasate-Mondragón, Spain

Introduction

Along the evolutionary path from single cells to multicellular organisms with a central nervous system, there are species of intermediate complexity that possess many autonomous cells programmed to respond to environmental stimuli. One of the more striking responses is phototaxis[1], in which motile organisms adjust their locomotory path according to the incident light in a finely controlled manner, either towards or against the light source (positive or negative phototaxis). Inspired by them we developed an inanimate/chemical system in which an organic droplet is self-propelled in response to a photo-stimulus. The centimetre-scale directional movement of the organic droplet on the aqueous solution is powered by the combination of photo-induced pH change and surface tension effects.






Photo-controlled pH

Dissolution of SP-SO₃H in water leads to dissociation of the sulfonic acid and ring opening of the SP to form an equilibrium mixture of yellow MCH⁺-SO₃⁻ (predominant) and red MC-SO₃⁻ (pH≈5). Upon white light irradiation, the equilibrium shifts as the open forms of the molecule (MCH⁺-SO₃⁻ and MC-SO₃⁻) are predominantly converted to SP-SO₃⁻ resulting in an increase in H⁺ conc. (pH≈3.4).




Phototactic droplets


The principle of phototaxis offers new possibilities to employ light as an external stimulus for controlling droplet motion at the liquid-air interface. This new concept is generic and can be applied to various organic/ aqueous systems, making photo-manipulation of microliter-size droplets relatively easy to implement. Here, depending on the intensity of the light source, movement of droplets with speeds in the range of 100 to 4000 $\mu\text{m s}^{-1}$ can be achieved.



Powered by Light



(A) Droplet is stable in the initial pH conditions (pH≈5, MC-H⁺ solution). (B) Upon irradiation with white light, MC-H⁺ reverts to SP, causing the generation of free H⁺ (pH≈3.4). (C) Under these acidic conditions, the DA⁻ ion at boundary region becomes protonated, and spontaneously migrates back into the bulk droplet region. Simultaneously, to preserve charge neutrality, the CI-H⁺ ion (surfactant) migrates into the aqueous phase; as soon as this happens, the surface tension of the solution dramatically drops, causing the droplet to spontaneously move to the opposite direction of the light source. (D) The droplet stops in the region of high surface tension, where it encounters the initial conditions (MC-H⁺, pH≈5).



[1] D. B. Weibel, P. Garstecki, D. Ryan, W. R. DiLuzio, M. Mayer, J. E. Seto, G. M. Whitesides, Proc. Natl. Acad. Sci. U.S.A., 2005, 102, 11963.

Acknowledgements : Science Foundation Ireland under the INSIGHT initiative, grant SFI/12/RC/2289 (INSIGHT), Australian Research Council and EU IRSES program.