

(I). Financial support from the Polish State Committee for Scientific Research (project 2P303 123 07) is gratefully acknowledged.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: KA1293). Services for accessing these data are described at the back of the journal.

References

- Allen, F. H., Bellard, S., Brice, M. D., Cartwright, B. A., Doubleday, A., Higgs, H., Hummelink, T., Hummelink-Peters, B. G., Kennard, O., Motherwell, W. D. S., Rodgers, J. R. & Watson, D. G. (1979). *Acta Cryst.* **B35**, 2331–2339.
- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1992). *International Tables for Crystallography*, Vol. C, edited by A. J. C. Wilson, pp. 691–705. Dordrecht: Kluwer Academic Publishers.
- Bertolasi, V., Gilli, P., Ferretti, V. & Gilli, G. (1994). *Acta Cryst.* **B50**, 617–625.
- Bertolasi, V., Nanni, P., Gilli, P., Ferretti, V., Gilli, G., Issa, Y. M. & Sherif, O. E. (1994). *New J. Chem.* **18**, 251–261.
- Bondi, A. (1964). *J. Phys. Chem.* **68**, 441–451.
- Dal Colle, M., Bertolasi, V., de Palo, M., Distefano, G., Jones, D., Modelli, A. & Olivato, P. R. (1995). *J. Phys. Chem.* **99**, 15011–15017.
- Distefano, G., Dal Colle, M., de Palo, M., Jones, D., Bombieri, G., Del Pra, A., Olivato, P. R. & Mondino, M. G. (1996). *J. Chem. Soc. Perkin Trans. 2*, pp. 1661–1669.
- Etter, M. C. (1990). *Acta Cryst.* **B46**, 256–262.
- Gómez-Sánchez, A., de Garcia Martin, G. & Borrachero, P. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. 301–306.
- Hansch, C. & Leo, A. (1979). *Substituents Constants for Correlation Analysis in Chemistry and Biology*, p. 131. New York: Wiley.
- Hassall, K. A. (1982). *The Chemistry of Pesticides. Their Metabolism, Mode of Action and Uses in Crop Protection*, p. 176. Weinheim: Verlag Chemie.
- Juaristi, E. & Cuevas, G. (1992). *Tetrahedron*, **48**, 5019–5087.
- Kirby, A. J. (1983). *The Anomeric Effect and Related Stereoelectronic Effects at Oxygen*, p. 32. Berlin: Springer.
- Mandell, G. L. & Sande, M. A. (1985). *The Pharmacological Basis of Therapeutics*, edited by A. G. Gilman, L. S. Goodman, T. W. Rall & F. Murad, pp. 1212–1213. New York: MacMillan.
- March, J. (1992). *Advanced Organic Chemistry. Mechanisms and Structure*, p. 51. New York: Wiley.
- Sheldrick, G. M. (1985). *SHELXS86. Program for the Solution of Crystal Structures*. University of Göttingen, Germany.
- Sheldrick, G. M. (1997). *SHELXL97. Program for the Refinement of Crystal Structures*. University of Göttingen, Germany.
- Siemens (1989). *P3 Diffractometer Control Program*. Version 2.0. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Siemens (1990). *XP. Interactive Molecular Graphics Program*. Version 4.1. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Siemens (1991). *XDISK. Data Reduction Program*. Version 4.20. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- TRIPOS Associates Inc. (1996). *SYBYL. Molecular Modeling Software*. Version 6.3. TRIPOS Associates Inc., 1699 S. Hanley Rd. Suite 303, St Louis, Missouri 63144–2913, USA.
- Zakrzewski, A. (1996). Private communication.

Acta Cryst. (1999). **C55**, 472–474

Intermolecular C—H···N and C—H···O interactions in (2*S*,4*S*,5*R*)-(–)-3,4-dimethyl-5-phenyl-2-(1,3-thiazol-2-yl)-1,3-oxazolidine

LAVELLE M. FITZSIMONS AND JOHN F. GALLAGHER

School of Chemical Sciences, Dublin City University, Dublin 9, Ireland. E-mail: gallagherjfg@dcu.ie


(Received 29 September 1998; accepted 28 October 1998)

Abstract

The title compound, C₁₄H₁₆N₂OS, prepared from (1*R*,2*S*)-(–)-ephedrine, contains the oxazolidine ring in an envelope conformation, with the nitrogen atom 0.623 (2) Å from the plane of the other four oxazolidine-ring atoms. Intermolecular C—H···N and C—H···O interactions generate a two-dimensional hydrogen-bonded network, with shortest C···N and C···O distances of 3.403 (3) and 3.463 (2) Å, respectively.

Comment

Amino acid derivatives continue to be an important class of chiral compounds with applications in asymmetric synthesis and catalysis. The title compound, (I), a derivative of (1*R*,2*S*)-(–)-ephedrine, (II), is of current interest as a chiral auxiliary and as a new potentially bidentate ligand with an O,N or O,S donor-atom set for use in coordination chemistry.



Bond lengths and angles are unexceptional and in accord with anticipated values (Orpen *et al.*, 1994). The absolute structure can be deduced from (II) and by analysis of our X-ray data. The oxazolidine ring adopts an envelope conformation with N3 0.623 (2) Å from the (O1,C2,C4,C5) plane, which is at angles of 74.18 (6) and 86.99 (6)° to the phenyl and thiazole rings, respectively. The phenyl ring is oriented at an angle of 41.15 (8)° to the thiazolyl group. The molecular geometry in (I) compares with (2*S*,4*S*,5*R*)-(–)-2-(1*H*-imidazol-2-yl)-3,4-dimethyl-5-phenyl-1,3-oxazolidine, (III) (Gallagher

et al., 1998), and a related *p*-bromophenyl derivative (Just *et al.*, 1983). Examination of the structure with PLATON (Spek, 1997a) showed that there were no solvent-accessible voids in the crystal lattice.




Fig. 1. A view of (I) with the atomic numbering scheme. Displacement ellipsoids are drawn at the 30% probability level.




Fig. 2. A view of the intermolecular interactions in the crystal structure of (I).

There is an intermolecular $Csp^3-H \cdots O$ interaction to O1 and a $C(\text{thiazole})-H \cdots N(\text{thiazole})$ interaction to N7, which generates a two-dimensional hydrogen-bonded network, as depicted in Fig. 2, with details in Table 2. The closest $H \cdots S$ distance is the intramolecular $H12 \cdots S1$ of 3.00 Å, which is equal to the sum of the van der Waals radii (Bondi, 1964). The crystal structure of (I) contrasts with that of (III) where $N-H \cdots N$ hydrogen bonds, $Csp^3-H \cdots \pi(C=C, \text{imidazole})$, $C(\text{arene})-H \cdots \pi(\text{arene})$ and $C(\text{arene})-H \cdots O$ interactions generate a three-dimensional network (Gallagher *et al.*, 1998). The difference in hydrogen-bonding abilities when comparing S(thiazolyl) in (I) and N-H(imidazolyl) in (III) accounts for the lack of strong hydrogen-bond formation and the presence of weak $C-H \cdots N$ and $C-H \cdots O$ hydrogen bonds in (I). Further comparative studies are in progress on related 1,3-oxazolidine systems.

Experimental

The compound was prepared by refluxing thiazole-2-carboxaldehyde (500 mg, 4.42 mmol) and (1*R*,2*S*)-(-)-ephedrine (730 mg, 4.42 mmol) in 20 cm³ of acetonitrile for 4 h. On cooling, the product was filtered and recrystallized from ethanol, yield 0.98 g (85%), m.p. 354–355 K. IR ν_{max} (KBr) cm⁻¹: 3097 (*m*), 2800 (*m*), 2377 (*m*), 1455 (*s*), 1340 (*s*), 1059 (*s*). $[\alpha]_D^{20} = -75^\circ$ ($c = 1.0$, ethanol). ¹H NMR (400 MHz) (δ , CDCl₃) p.p.m.: 0.78 (*d*, 3H, CCH₃), 2.39 (*s*, 3H, NCH₃), 3.06 (*m*, 1H, MeCH), 5.12 (*s*, 1H, OCH), 5.20 (*d*, 1H, PhCH), 7.30–7.46 (*m*, 5H, C₆H₅), 7.46 (*d*, 1H, thiazole SCH), 7.83 (*d*, 1H, thiazole NCH).

Crystal data

C₁₄H₁₆N₂OS
 $M_r = 260.35$
 Orthorhombic
 $P2_12_12_1$
 $a = 10.2981(8) \text{ \AA}$
 $b = 10.8896(9) \text{ \AA}$
 $c = 12.1733(7) \text{ \AA}$
 $V = 1365.14(17) \text{ \AA}^3$
 $Z = 4$
 $D_x = 1.267 \text{ Mg m}^{-3}$
 D_m not measured

Mo $K\alpha$ radiation
 $\lambda = 0.7107 \text{ \AA}$
 Cell parameters from 25 reflections
 $\theta = 9.73\text{--}19.75^\circ$
 $\mu = 0.227 \text{ mm}^{-1}$
 $T = 294(1) \text{ K}$
 Block
 $0.30 \times 0.30 \times 0.29 \text{ mm}$
 Colourless

Data collection

Enraf–Nonius CAD-4 diffractometer
 ω -2 θ scans
 Absorption correction: none
 3406 measured reflections
 3118 independent reflections
 2613 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.012$
 $\theta_{\text{max}} = 27.4^\circ$
 $h = -13 \rightarrow 13$
 $k = 0 \rightarrow 14$
 $l = 0 \rightarrow 15$
 3 standard reflections
 frequency: 120 min
 intensity variation: 0.5%

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.091$
 $S = 1.024$
 3118 reflections
 164 parameters
 H-atom parameters
 constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0481P)^2 + 0.1202P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.15 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.21 \text{ e } \text{\AA}^{-3}$
 Extinction correction:
SHELXL97 (Sheldrick, 1997)
 Extinction coefficient:
 0.0120 (18)
 Scattering factors from
International Tables for Crystallography (Vol. C)
 Absolute structure: Flack (1983)
 Flack parameter = -0.01 (8)

Table 1. Selected geometric parameters (\AA , $^\circ$)

S1—C6	1.7250 (18)	C2—C6	1.483 (2)
S1—C9	1.703 (2)	N7—C6	1.289 (2)
O1—C2	1.414 (2)	N7—C8	1.373 (3)
O1—C5	1.445 (2)	C8—C9	1.337 (3)
C6—S1—C9	89.54 (10)	O1—C5—C11	110.17 (15)
C2—O1—C5	108.36 (14)	S1—C6—N7	114.31 (14)
O1—C2—N3	104.45 (14)	S1—C6—C2	121.20 (12)
O1—C2—C6	109.74 (14)	S1—C9—C8	109.48 (16)
O1—C5—C4	104.16 (15)		
O1—C2—C6—S1	81.84 (16)	O1—C5—C11—C12	-38.7 (2)
O1—C2—C6—N7	-95.7 (2)	O1—C5—C11—C16	141.50 (17)

Table 2. Hydrogen-bonding geometry (\AA , $^\circ$)

D—H...A	D—H	H...A	D...A	D—H...A
C9—H9...N7	0.93	2.73	3.403 (3)	130
C41—H41C...O1 ^b	0.96	2.62	3.463 (2)	147

Symmetry codes: (i) $\frac{1}{2} + x, \frac{1}{2} - y, 2 - z$; (ii) $\frac{1}{2} + x, \frac{1}{2} - y, 2 - z$.

Molecule (I) is chiral and a complete 'Friedel' data set ($\pm h, +k, +l$) was collected and the absolute structure determined [Flack parameter -0.01 (8)] by our X-ray analysis. The known absolute configuration of the (1*R*,2*S*)-(-)-ephedrine starting material, (II), is retained in the three chiral centres (2*S*,4*S*,5*R*) of the oxazolidine ring in (I). H atoms were allowed for as riding atoms with C—H in the range of 0.93 to 0.98 \AA .

Data collection: *CAD-4-PC Software* (Enraf-Nonius, 1992). Cell refinement: *SET4* and *CELDIM* in *CAD-4-PC Software*. Data reduction: *DATRD2* in *NRCVAX96* (Gabe *et al.*, 1989). Program(s) used to solve structure: *SHELXS97* (Sheldrick, 1990). Program(s) used to refine structure: *NRCVAX96* and *SHELXL97* (Sheldrick, 1997). Molecular graphics: *NRCVAX96*, *ORTEP* (Johnson, 1976), *PLATON* (Spek, 1997*a*) and *PLUTON* (Spek 1997*b*). Software used to prepare material for publication: *NRCVAX96*, *SHELXL97* and *WordPerfect* macro *PRPCIF97* (Ferguson, 1997).

JFG thanks Dublin City University and Forbairt for funding of a research visit to the University of Guelph in 1998 and especially Professor George Ferguson for use of his diffractometer and computer system.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: BM1296). Services for accessing these data are described at the back of the journal.

References

- Bondi, A. (1964). *J. Phys. Chem.* **68**, 441–451.
 Enraf-Nonius (1992). *CAD-4-PC Software*. Version 1.1. Enraf-Nonius, Delft, The Netherlands.
 Ferguson, G. (1997). *PRPCIF97. A WordPerfect-5.1 Macro to Merge and Polish CIF Format Files from NRCVAX and SHELXL97 Programs*. University of Guelph, Canada.
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
 Gabe, E. J., Le Page, Y., Charland, J.-P., Lee, F. L. & White, P. S. (1989). *J. Appl. Cryst.* **22**, 384–387.
 Gallagher, J. F., Briody, J. M. & Cantwell, B. P. (1998). *Acta Cryst.* **C54**, 1331–1335.
 Johnson, C. K. (1976). *ORTEP*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
 Just, G., Potvin, P., Uggowitzer, P. & Bird, P. (1983). *J. Org. Chem.* **48**, 2923–2924.
 Orpen, A. G., Brammer, L., Allen, F. H., Kennard, O., Watson, D. G. & Taylor, R. (1994). *Structure Correlation*, Vol. 2, edited by H.-B. Bürgi & J. D. Dunitz, Appendix A. Weinheim: VCH Publishers.
 Sheldrick, G. M. (1990). *Acta Cryst.* **A46**, 467–473.
 Sheldrick, G. M. (1997). *SHELXL97. Program for the Refinement of Crystal Structures*. University of Göttingen, Germany.
 Spek, A. L. (1997*a*). *PLATON. Molecular Geometry Program*. Version of May 1997. University of Utrecht, The Netherlands.
 Spek, A. L. (1997*b*). *PLUTON. Molecular Graphics Program*. Version of May 1997. University of Utrecht, The Netherlands.

Acta Cryst. (1999). **C55**, 474–476

Bis(*m*-anisidinium) dihydrogénéodiphosphate

MOHAMED OULD ABDELLAHI, FATMA BEN AMOR, AHMED DRISS ET TAHAR JOUINI

Département de Chimie, Faculté des Sciences, 1060 Campus Universitaire, Tunis, Tunisie. E-mail: tahar.jouini@fst.rnu.tn

(Reçu le 14 mars 1997, accepté le 7 septembre 1998)

Abstract

The structure consists of an alternate stacking of $(\text{H}_2\text{P}_2\text{O}_7)^{2-}$ anions and $(1,3\text{-CH}_3\text{OC}_6\text{H}_4\text{NH}_3)^+$ cations held together by N—H...O intermolecular hydrogen bonds.

Commentaire

L'arrangement moléculaire de cette structure, (I), est constitué de couches d'anions phosphates $(\text{H}_2\text{P}_2\text{O}_7)^{2-}$

